


This article was downloaded by: [Ruud H.J. Hornsveld]

On: 18 August 2014, At: 09:58

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH,
UK



The Journal of Forensic Psychiatry & Psychology

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/rjfp20>

Behavior on the ward of personality-disordered inpatients and chronically psychotic inpatients during a three-year stay in a Dutch forensic psychiatric hospital

Ruud H.J. Hornsveld^a, Floris W. Kraaimaat^b, Samantha Bouwmeester^c, Machiel A. Polak^d & Almar J. Zwets^{cd}

^a Erasmus University Medical Centre, Rotterdam, The Netherlands

^b Radboud University, Nijmegen, The Netherlands

^c Institute for Psychology, Erasmus University, Rotterdam, The Netherlands

^d FPC de Kijvelanden, Poortugaal, The Netherlands

Published online: 14 Aug 2014.

To cite this article: Ruud H.J. Hornsveld, Floris W. Kraaimaat, Samantha Bouwmeester, Machiel A. Polak & Almar J. Zwets (2014): Behavior on the ward of personality-disordered inpatients and chronically psychotic inpatients during a three-year stay in a Dutch forensic psychiatric hospital, *The Journal of Forensic Psychiatry & Psychology*, DOI: [10.1080/14789949.2014.947308](https://doi.org/10.1080/14789949.2014.947308)

To link to this article: <http://dx.doi.org/10.1080/14789949.2014.947308>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no

representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Behavior on the ward of personality-disordered inpatients and chronically psychotic inpatients during a three-year stay in a Dutch forensic psychiatric hospital

Ruud H.J. Hornsveld^{a*}, Floris W. Kraaimaat^b, Samantha Bouwmeester^c,
Machiel A. Polak^d and Almar J. Zwets^{c,d}

^aErasmus University Medical Centre, Rotterdam, The Netherlands; ^bRadboud University, Nijmegen, The Netherlands; ^cInstitute for Psychology, Erasmus University, Rotterdam, The Netherlands; ^dFPC de Kijvelanden, Poortugaal, The Netherlands

(Received 6 December 2013; accepted 16 July 2014)

To gain insight into the behavior of a group of personality-disordered patients and a group of chronically psychotic patients during their stay in a Dutch forensic psychiatric hospital, data on these patients were collected biannually for seven years. Three aspects of the patients' behavior were examined: the prediction of institutional behavior shortly after admission, changes in the patients' behavior on the ward during their stay in hospital, and the prediction of these changes. In the personality-disordered patients, observed irritation/anger and aggressive behavior on the ward turned out to be positively related to psychopathy, the PCL-R lifestyle and antisocial facets, and the neuroticism domain. A positive relationship was also found between aggressive behavior on the ward and trait anger. In the chronically psychotic patients, a positive relationship was found between irritation/anger and the PCL-R interpersonal and lifestyle facet. During a stay of three years, the aggressive behavior of both patient subgroups, which was already low at the start, did not decrease further, but their prosocial behavior increased. In the personality-disordered patients, relatively high scores on the antisocial facet of the PCL-R indicated an increase in prosocial behavior, whereas in the chronically psychotic patients no relationship was found between any PCL-R facet and behavior change. Effect studies on treatment programs for forensic psychiatric inpatients have to contend with the problem of a low base rate of institutional aggression. Therefore, we advise that such studies focus not only on a decrease in negative behaviors but also on an increase in positive behaviors.

Keywords: forensic psychiatry; personality-disordered inpatients; chronically psychotic inpatients; behavior on the ward

Introduction

To gain insight into the behavior of a group of personality-disordered patients and a group of chronically psychotic patients during their stay in a Dutch

*Corresponding author. Email: r.hornsveld@tiscali.nl

forensic psychiatric hospital, data on these patients were collected biannually and prospectively for seven years. Three aspects of the patients' behavior were examined: the prediction of institutional behavior shortly after admission, changes in the patients' behavior on the ward during a three-year stay in hospital, and the prediction of these changes.

During the past decade, a considerable number of studies devoted to the relationship between psychopathy and recidivism and to the relationship between psychopathy and institutional aggression have been published. For instance, Hildebrand, De Ruiter, and Nijman (2004) investigated the predictive validity of the Psychopathy Checklist-Revised (PCL-R; Hare, 1991) and its two factors in a group of 92 Dutch forensic psychiatric inpatients and found that factor 2 ('Chronically unstable and antisocial lifestyle') was the best predictor for the total number of institutional incidents, but not for physical violence on the ward. Guy, Edens, Anthony, and Douglas (2005) performed a meta-analysis of 273 effect sizes to investigate the associations between the PCL-R and institutional misconduct. Their results suggested that factor 2 was associated with misconduct more strongly than factor 1 ('Callous and remorseless use of others'), but for both factors, the associations with physical violence were smaller than the associations with misconduct. Taking 95 American, Canadian, and European studies altogether, Leistico, Salekin, DeCoster, and Rogers (2008) performed a meta-analysis on the relationship between the various editions of the PCL (Hare, 1980) and antisocial conduct, both institutional misbehavior and recidivism after discharge. The results indicated that higher scores on PCL total, factor 1, and factor 2 were moderately associated with increased antisocial conduct. However, effect sizes depended on variables such as country, race, gender, and institutional setting. For instance, the mean effect sizes of factor 2 were found to be slightly larger in samples of forensic psychiatric patients than in samples of detainees. Two years later, Yang, Wong, and Coid (2010) found more or less similar results in comparing the effect sizes of nine risk-assessment instruments from 28 original reports, namely that the predictive efficacy of the PCL-R for violence was attributable almost entirely to factor 2. In order to find out whether the combination of factor 1 and factor 2 would result in a better prediction of violence than only factor 2, Kennealy, Skeem, Walters, and Camp (2010) conducted a meta-analysis of 32 effect sizes. Again, factor 2 turned out to predict violence better than factor 1, but no interaction was found between the two factors. Although the former two studies supported the evidence for PCL-R factor 2 in being a predictor of future violence, it should be noted that the meta-analyses in these studies overlapped partly with the ones of the Leistico et al.'s (2008) study and no differentiation was made between forensic psychiatric patients and offenders, and between institutional and community violence.

After Hare's (2003) introduction of the PCL-R four-facet model, the emphasis shifted from factor 2 to facet 4 in studies on the predictive validity of this instrument (e.g. Walters, Knight, Grann, & Dahle, 2008). In 2010,

Walters and Heilbrun published a study on the relationship between facet 4 ('Antisocial') of the PCL-R and antisocial conduct in a sample of 216 male forensic patients with the PCL (Hare, 1980) and in a sample of 230 psychiatrically evaluated inmates with the PCL-R (Hare, 1991). They found, in accordance with the Walters et al.'s study (2008) that facet 4 or parcel G (items 'early behavioral problems' and 'poor behavioral control') consistently achieved incremental validity to the first three facets ('Interpersonal,' 'Affective,' and 'Lifestyle'). Although several authors (Hildebrand et al., 2004; Leistico et al., 2008; Walters & Heilbrun, 2010) investigated the predictive validity of the PCL-R in (among others) forensic psychiatric patients, no differentiation was made between patients with a major mental disorder on Axis I of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR; American Psychiatric Association, 2000) and those with a primary diagnosis on Axis II in diagnostically mixed samples. Furthermore, Vitacco et al. (2009) suggested that when investigating the utility of the PCL-R and its factors for the prediction of institutional aggression, a differentiation should be made between reactive and proactive aggression on the ward. In a group of 152 mainly psychotic forensic psychiatric inpatients, they showed that psychopathy (PCL:SV; Hart, Cox, & Hare, 1995) and its Interpersonal and Affective facets were robust predictors for proactive aggression, but not for reactive aggression. However, in explaining the various results of the studies, several authors have emphasized that the PCL-R has been designed as an instrument for assessing psychopathy and not for assessing the risk of future institutional or community violence. Therefore, they recommend that clinicians and legal decision-makers consider risk and protective factors beyond psychopathy when attempting to predict future behaviors (e.g. Edens, Campbell, & Weir, 2007; Leistico et al., 2008; Singh, Grann, & Fazel, 2011).

Contrary to numerous publications about the relationship between psychopathy and antisocial behavior, the number of publications about the relationship between personality traits as measured by self-report questionnaires and institutional behavior is very limited. Until now, no study has been carried out on the relationship between the Big Five personality domains (Costa & McCrae, 1989) and institutional behavior. The only study about the relationship between anger as a trait and institutional behavior was performed by Cornell, Peterson, and Richards (1999), in a group of 65 incarcerated adolescents. Trait anger, as measured by the State-Trait Anger Expression Inventory (STAXI; Spielberger, 1988) appeared to be significantly correlated with physical and with verbal aggression as rated by the staff.

Studies on the effects of institutional treatment programs for forensic psychiatric patients have yielded divergent results. Recently, McGuire (2013) concluded that cognitive-behavioral treatment programs for violent offenders have positive effects on recidivism, but studies on the behavior change in forensic psychiatric patients during their stay in hospital usually show less convincing results. For instance, Belfrage and Douglas (2002) studied 70 Swedish,

mostly psychotic patients with the 20-item Historical/Clinical/Risk Management-20 (Webster, Douglas, Eaves, & Hart, 1997). After an 18-month stay, patients scored significantly lower on the Clinical subscale and the Risk Management subscale. De Jonge, Nijman, and Lammers (2009) had similar results with a study of 158 Dutch, mostly personality-disordered forensic psychiatric patients. After 25 months, patients scored significantly lower on both the Clinical factor and the Future factor of the History/Clinical/Future 30 (Ministry of Security and Justice, 2002), but effect sizes were small. Nijman, De Kruyk, and Van Nieuwenhuizen (2004) studied the course of behavior by means of the Rehabilitation Evaluation Hall and Baker (Baker & Hall, 1988) in a group of 128 Dutch, mainly personality-disordered forensic psychiatric inpatients. During hospitalization, social activity improved significantly, but improvements in self-care and verbal skills were not significant. Chakhssi, De Ruiter, and Bernstein (2010) studied a group of 74 Dutch, personality-disordered inpatients for a period of 20 months. Using the Behavioral Status Index (BSI; Reed, Woods, & Robinson, 2000), they found that most patients 'improved' to a limited extent on the BSI, but the psychopaths (22%) 'deteriorated' during the study period, whereas none of the non-psychopaths did. Finally, Hildebrand and De Ruiter (2012) used self-report questionnaires, the Rorschach Inkblot Method (Rorschach, 1921/1942), staff ratings, and objective measures for treatment compliance for their investigation in a group of 87 Dutch, forensic psychiatric inpatients. The group of patients as a whole did not improve on most of the indicators of dynamic risk after 20 months of treatment. A few researchers (Belfrage & Douglas, 2002; Hildebrand & De Ruiter, 2012) suggest that the limited change in the behavior of forensic psychiatric patients during their stay in a hospital might be due to treatment programs, which deserve review or alteration. However, differences in results between the studies about institutional behavior and the studies as summarized by McGuire (2013) might also be explained by differences in outcome measures, namely reduction of dynamic criminogenic needs versus reduction of recidivism rate.

In the current study, it was hypothesized (1) that psychopathy as assessed by the PCL-R was related to institutional aggression in forensic psychiatric inpatients with a personality disorder, as well as in inpatients with a chronically psychotic disorder, and that this relationship involved specifically facet 4 of the PCL-R; (2) that in each of these patient groups, the hospital stay would result in a decrease of aggressive behavior and, in an increase of prosocial behavior; and (3) that these behavior changes were related to psychopathy in both inpatient groups.

Methods

Participants

In the Netherlands, offenders who have committed a serious violent crime that is punishable with a maximum imprisonment of more than four years

(e.g. murder, manslaughter, aggravated assault, or rape) can be detained under hospital order ('TBS order'). This concerns offenders who, based on an extensive psychiatric and/or psychological evaluation at a special assessment center of the Ministry of Security and Justice, are judged to have diminished responsibility for the offense they committed (Van Marle, 2002). TBS involves involuntary admission to a specialized maximum-security forensic psychiatric hospital with obligatory treatment programs that should result in a decrease of recidivism risk to an 'acceptable level for society.' The Dutch Ministry of Security and Justice makes a distinction between patients with a 'personality disorder' (about 75% of the population) and patients with a 'chronically psychotic disorder' (De Beurs & Barendregt, 2008).

Behavior on the ward was assessed prospectively and biannually (in the months of January and July) as part of a standard procedure between January 2003 and January 2011 by means of the Observation Scale for Aggressive Behavior (OSAB; Hornsveld, Nijman, Hollin, & Kraaimaat, 2007). During this period, the number of assessed patients gradually increased because of new admissions. Ultimately, data on 253 patients could be collected with one measurement, on 248 patients with two measurements, on 236 patients with three measurements, and so on (Table 1). The mean age of the 253 patients with one measurement (shortly after admission) was 37.46 years (SD = 10.42, range: 19–69 years). The 159 patients of the group classified as personality disordered had a mean age of 38.59 years (SD = 10.67, range: 19–65 years) at admission, and the 94 classified as chronically psychotic patients had a mean age of 35.54 years (SD = 9.73, range: 22–69 years).

Because the average stay of Dutch forensic psychiatric inpatients has increased dramatically in recent years, reaching 9.8 years in 2010 (Nachtegaal,

Table 1. Number of patients with number of measurement moments, admitted between January 2003 and January 2008.

Measurement moments	Total group of patients		Personality-disordered patients		Chronically psychotic patients	
	<i>N</i>	Age	<i>n</i>	Age	<i>n</i>	Age
1	253	37.46 (10.42)	159	38.59 (10.67)	94	35.54 (9.73)
2	248	37.40 (10.44)	157	38.58 (10.68)	91	35.37 (9.75)
3	236	37.17 (10.41)	148	38.38 (10.66)	88	35.15 (9.69)
4	213	37.20 (10.34)	134	38.69 (10.88)	79	34.67 (8.86)
5	178	36.97 (10.46)	108	38.69 (11.14)	70	34.30 (8.74)
6	146	37.03 (10.17)	86	38.99 (10.77)	60	34.22 (8.58)
7	115	36.97 (10.27)	70	38.79 (10.75)	45	34.13 (8.85)
8	84	36.68 (9.57)	54	39.09 (10.22)	30	32.33 (6.40)
9	70	37.03 (9.66)	47	39.11 (10.29)	23	32.78 (6.58)
10	48	35.75 (8.91)	30	37.50 (9.90)	18	32.83 (6.17)
11	24	36.50 (8.40)	16	37.19 (9.49)	8	35.13 (5.96)

Van der Horst, & Schönberger, 2011), we chose a period of three years for an analysis of behavior on the ward, a compromise between a relatively long period of stay and a sufficient number of patients. In total, 115 patients could be assessed through the OSAB on seven measurement moments from admission onwards. At admission, the 70 personality-disordered patients of this group had a mean age of 38.79 years (SD = 10.75, range 19–62 years), and the 45 chronically psychotic patients had a mean age of 34.13 years (SD = 8.85, range 22–68 years). Some of these 115 patients were admitted in the second half of 2002, were measured for the first time in January 2003, and were measured for the seventh time in January 2006. Other patients were admitted in the first half of 2003, were measured for the first time in July 2003, and were measured for the seventh time in July 2006. The last of these 115 patients were admitted in the second half of 2007, were measured for the first time in January 2008, and were measured for the seventh time in January 2011. Missing scores (e.g. during the fifth measurement) were added by calculating the average score of the two adjacent scores (e.g. during the fourth and sixth measurements). The percentage of missing scores in this part of the data-set was 8%.

In the period between January 2003 and January 2008, 204 patients were admitted to the hospital. Data on the remaining 89 patients were incomplete because of several reasons. For 45 patients (group 2), data from less than five measurements could be collected, although they had stayed in the institution for more than three years. Others stayed for less than three years in hospital during the investigated period, because they were discharged (7 patients, group 3), transferred to another institution for reassessment (20 patients, group 4), or had to continue their care in a long-stay hospital elsewhere (6 patients, group 5). Furthermore, 11 patients (group 6) were admitted only for a very short time, for instance, in case of a crisis intervention (Table 2). Groups 2–6 did not differ in PCL-R total score from group 1, which included the 115 patients with seven measurements.

Measures

The *Observation Scale for Aggressive Behavior* (OSAB; Hornsveld et al., 2007) measures behavior on the ward. The scale comprises 40 items spread over the subscales Irritation/Anger, Anxiety/Gloominess, Aggressive Behavior, Prosocial Behavior, Antecedent, and Sanction. The staff judges the behavior of the inpatients in the preceding week on a four-point scale, with 1 = ‘no,’ 2 = ‘seldom,’ 3 = ‘occasionally,’ and 4 = ‘frequently.’ In this study, we focused on the following subscales: Irritation/Anger (e.g. ‘Irritated’), Aggressive Behavior (e.g. ‘Threats toward staff’), and Prosocial Behavior (e.g. ‘Gives his opinion adequately’). By far, most of these aggressive and prosocial behaviors was exhibited on the ward in relation to staff members. Aggressive behavior towards fellow patients probably took place in the absence of staff

Table 2. Mean scores and standard deviations for age, PCL-R, and OSAB in all patients who were admitted in the hospital from January 2003 to January 2008.

Number	Group	Number of patients <i>n</i>	Percent- tage %	Age <i>M (SD)</i>	PCL-R Total <i>M (SD)</i>	OSAB		
						Irritation/ anger <i>M (SD)</i>	Aggressive behavior <i>M (SD)</i>	Prosocial behavior <i>M (SD)</i>
1	Three-year stay in hospital with seven measurements	115	56.4	36.97 (10.27)	20.49 (7.97)	10.19 (3.62)	14.97 (5.66)	17.43 (8.60)
2	Three-year stay in hospital with less than five measurements	45	22.1	38.62 (11.95)	18.00 (7.87)	11.32 (3.76)	15.92 (4.60)	29.11 (7.73)
3	End of TBS or on leave	7	3.4	41.29 (11.94)	17.00 (8.25)	10.14 (2.48)	14.14 (2.48)	35.29 (4.50)
4	Transfer or reselection	20	9.8	36.95 (8.89)	22.63 (7.86)	11.63 (3.67)	16.16 (5.23)	27.89 (7.64)
5	Long-stay hospital	6	2.9	55.00 (8.46)	23.40 (7.57)	13.20 (2.49)	18.40 (2.61)	30.80 (5.98)
6	Very short stay	11	5.4	35.89 (6.31)	25.67 (9.35)	11.22 (2.59)	15.67 (3.20)	27.44 (6.00)

Note: PCL-R = psychopathy checklist-revised; OSAB = observation scale for aggressive behavior.

members. In a sample of 220 violent forensic psychiatric inpatients, the internal consistency (Cronbach's α) was .82, .79, and .93, inter-rater reliability was .79, .81, and .70, and test-retest reliability was .59, .57, and .76, for the subscales Irritation/Anger, Aggressive Behavior, and Prosocial Behavior, respectively (Hornsveld et al., 2007). Convergent validity was supported through significant positive correlations with subscales of the Forensic Inpatient Observation Scale (Timmerman, Vastenburg, & Emmelkamp, 2001).

The PCL-R (Hare, 1991; Dutch version: Vertommen, Verheul, De Ruiter, & Hildebrand, 2002) was employed to measure psychopathy. The checklist consists of 20 items, which have to be rated on a three-point scale, with 0 = 'does not apply,' 1 = 'applies to some extent,' and 2 = 'applies.' Vertommen et al. (2002) found support for the reliability of the Dutch version of the PCL-R in a group of 1192 inmates. The internal consistency (Cronbach's α) was .87. Tentative support for convergent validity was found in a subgroup of 98 forensic psychiatric inpatients, as there were modest but meaningful correlations with self-report questionnaires such as the MMPI-2 (Dutch version: Sloore, Derksen, Hellenbosch, & De Mey, 1993). In the present study, we used the total score as well as the four-factor structure as proposed by Hare and Neumann (2006), which implies the following reliable facets: Interpersonal, Affective, Lifestyle, and Antisocial. The PCL-R scores of a sample of 41 patients was assessed independently by two trained psychologists. Intra-class correlation coefficient for the PCL-R total score was .81, and for the four facets .66, .64, .58, and .86 successively. The coefficient of the PCL-R total score was regarded as excellent, the coefficient of facet 2 as fair, the coefficients of facet 1 and facet 3 as good, and the coefficient of facet 4 as excellent (Cicchetti, 1994).

The *NEO Five-Factor Inventory* (NEO-FFI; Costa & McCrae, 1992; Dutch version: Hoekstra, Ormel, & De Fruyt, 1996) includes 60 items and measures the Big Five personality domains of neuroticism, extraversion, openness, agreeableness, and conscientiousness. Participants score the items in the NEO-FFI on a five-point Likert scale ranging from 'entirely disagree' to 'entirely agree.' In the present study, we were only interested in the neuroticism (e.g. 'I seldom feel lonely or sad.') and agreeableness (e.g. 'Some people find me selfish and egotistic.') scales, because these traits are considered relevant in the context of aggression (Hornsveld, Nijman, & Kraaimaat, 2008). In a Dutch sample of 356 non-clinical adults, the internal consistency (Cronbach's α) of the two subscales was .82 and .69; and in a subgroup of 135 adults, the test-retest reliability after 6 months was .82 and .75, respectively (Hoekstra et al., 1996).

The Trait Anger subscale of the Spielberger (1980) *State-Trait Anger Scale* (STAS; Van der Ploeg, Defares, & Spielberger, 1982), which consists of 10 items, was used as a concurrent measure of the general disposition to anger. Participants rate each item about how they generally feel (e.g. 'I am quick tempered.') by using a four-point Likert scale: 1 = 'almost never,' 2 = 'sometimes,' 3 = 'often,' and 4 = 'almost always.' In a group of 150 Dutch male university

students, Van der Ploeg et al. (1982) found that the internal consistency (Cronbach's α) of the trait anger scale was .78, and a test-retest reliability of .78 was documented in a subgroup of 70 students. The convergent validity of the trait anger scale also appeared to be satisfactory (Van der Ploeg et al., 1982).

Procedure

The study was approved by the Dutch Review Committee for Patient-linked Research in Arnhem, the Netherlands, and by the Scientific Research and Documentation Center of the Dutch Ministry of Security and Justice.

Group leaders were trained in the scoring of the OSAB, and scored this instrument biannually in the months of January and July. Certified clinical psychologists assessed the PCL-R scores based on file study. Such files comprised detailed information about life history, committed offenses, and elaborated evaluations from psychiatrists and/or psychologists. These reports were drawn up at a special forensic assessment center, in which the offender had to stay for observation by order of the court. The psychologists who scored the PCL-R were not informed about the OSAB scores.

Patients completed the questionnaires individually under supervision of an experienced research assistant and received a fee of € 7 in return for their participation. Not all patients completed the questionnaires, because participation in the study was on a voluntarily basis.

Setting

The current study was conducted at FPC de Kijvelanden in Poortugaal, the Netherlands. The patient-staff ratio was 1-1.8. All patients resided on high-security wards for seven to eleven patients, where specifically educated group leaders offered them milieu therapy. During a period of about four months after admission, the patients' behaviors were observed on the ward, psychiatric and psychological evaluations were carried out, and a treatment plan was finally established. Depending on their dynamic criminogenic needs, patients followed cognitive-behavioral treatment programs, focusing on (sexual) violence, addiction, or chronically psychotic disorders. When indicated, they also had to follow individual therapy, creative art therapy (e.g. drama, art, music, or psychomotor therapy), general education, occupational training, or sports. Psychopharmacological therapy was applied to all chronically psychotic patients and to the personality-disordered patients for whom it was indicated and who did not refuse medication.

Statistics

For the two groups separately, Pearson correlation coefficients were calculated for the relationship between scores on the PCL-R, NEO-FFI, or STAS on the

one hand, and the three OSAB subscales shortly after admission on the other hand (two-tailed, $p < .05$). In addition, stepwise multiple regression analyses were performed to explore the relative contribution of the four PCL-R facets to the three OSAB subscales in the two patient groups separately. To study mean linear changes in behavior at the ward as measured by the three OSAB subscales over time (seven measurements, biannually), three separate mixed analyses of variance were performed with groups as between-subject factor and the seven measurement moments as within-subject factor. In case of significant relationships, *post hoc* tests were performed. To explore the relationship between independent variables at time of admission and changes in behavior at the ward over a period of three years, Pearson correlation coefficients were calculated between the scores on PCL-R, NEO-FFI, and STAS at the time of admission and the residual gain scores for the three OSAB scales at measurement 7 (Kerlinger, 1975).

Results

Behavior on the ward shortly after admission

Table 3 shows the mean scores and standard deviations for all measures and 253 patients in each of the two samples.

Personality-disordered patients had a significantly higher total score on the PCL-R and its Interpersonal, Affective, and Lifestyle facets than the chronically psychotic patients. No significant difference between subgroups was found on the other measures, with the exception of the Irritation/Anger and Prosocial Behavior subscales of the OSAB, on which the personality-disordered patients scored significantly higher than the chronically psychotic patients.

Relationships between OSAB scores on the one hand and PCL-R, NEO-FFI, or STAS scores on the other hand were significant and in the expected direction for the personality-disordered patients but not for the chronically psychotic patients. In the personality-disordered group, the OSAB subscales Irritation/Anger and Aggressive Behavior correlated significantly positively with the total score on the PCL-R, the Lifestyle facet, the Antisocial facet, and the NEO-FFI domain Neuroticism. The OSAB subscale was also found to correlate significantly positively with the STAS. In the chronically psychotic group, the subscale Irritation/Anger correlated significantly positively with the Interpersonal and Lifestyle facet of the PCL-R. In the latter group, however, no significant relationships were found between the three OSAB subscales and the two NEO-FFI domains or the STAS (Table 4). Stepwise multiple regression analyses were conducted to evaluate which facets of the PCL-R might predict irritation/anger, aggressive behavior, or prosocial behavior shortly after admission. In the personality-disordered group, results indicated that the antisocial facet was significantly related to irritation/anger, $F(1, 157) = 14.42$,

Table 3. Mean scores and standard deviations for PCL-R, NEO-FFI, STAS, and OSAB in personality-disordered ($n = 159$) and chronically psychotic patients ($n = 94$).

Measure	Factor or subscale	Personality disordered		Chronically psychotic		Differences between subgroups		Cohen's d
		n	M (SD)	n	M (SD)	t		
Age		159	38.59 (10.67)	94	35.54 (9.73)		$t(251) = 2.26$ ($p = .024$)	.30
PCL-R	Psychopathy	159	22.25 (8.06)	94	17.96 (7.84)		$t(251) = 4.14$ ($p < .001$)	.54
	Interpersonal	159	3.57 (2.47)	94	1.80 (1.88)		$t(251) = 6.01$ ($p < .001$)	.81
	Affective	159	6.13 (1.72)	94	5.64 (1.84)		$t(251) = 2.08$ ($p = .039$)	.28
	Lifestyle	159	5.94 (2.76)	94	4.98 (2.90)		$t(251) = 2.62$ ($p = .009$)	.34
	Antisocial	159	5.11 (2.81)	94	4.68 (2.81)		$t(251) = 1.17$ ($p = .245$)	.15
NEO-FFI	Neuroticism	97	32.24 (8.49)	48	31.73 (7.92)		$t(143) = 0.35$ ($p = .729$)	.06
	Agreeableness	97	41.59 (5.23)	48	42.52 (4.93)		$t(143) = -1.03$ ($p = .305$)	-.18
STAS	Trait anger	92	17.91 (6.64)	47	15.85 (4.29)		$t(137) = 1.93$ ($p = .056$)	.38
OSAB	Irritation/anger	159	11.08 (3.42)	94	10.02 (3.84)		$t(251) = 2.28$ ($p = .024$)	.29
	Aggressive behavior	159	15.61 (5.05)	94	14.81 (5.33)		$t(251) = 1.20$ ($p = .233$)	.15
	Prosocial behavior	159	29.95 (7.80)	94	25.21 (7.82)		$t(251) = 4.67$ ($p < .001$)	.61

Note: PCL-R = psychopathy checklist-revised; NEO-FFI = five-factor inventory; STAS = state-trait anger scale; OSAB = observation scale for aggressive behavior.

Table 4. Correlations (Pearson's r) between scores on PCL-R, NEO-FFI, STAS, and OSAB for personality-disordered ($n = 159$) and chronically psychotic patients ($n = 94$), assessed shortly after admission (measurement 1).

Measure	Factors or subscales	N	Personality-disordered patients			OSAB			Chronically psychotic patients		
			Irritation/ Anger	Aggressive behavior	Prosocial behavior	n	Irritation/ anger	Aggressive behavior	Prosocial behavior	Irritation/ anger	Aggressive behavior
PCL-R	Psychopathy	159	.24**	.21**	.02	94	.17	.05	.08	.05	.08
	Interpersonal	159	.10	.07	.06	94	.19*	.06	.09	.06	.09
	Affective	159	.18*	.12	-.05	94	.08	-.01	.07	-.01	.07
	Lifestyle	159	.21**	.20**	-.00	94	.25**	.16	.03	.16	.03
NEO-FFI	Antisocial	159	.29**	.27**	.09	94	.05	-.03	.04	-.03	.04
	Neuroticism	97	.20*	.21*	-.11	48	.06	-.00	.16	-.00	.16
STAS	Agreeableness	97	-.10	-.16	.09	48	-.11	-.13	-.10	-.13	-.10
	Trait anger	92	.14	.21*	.02	47	.16	.18	.07	.18	.07

Note: PCL-R = psychopathy checklist-revised; NEO-FFI = five-factor inventory; STAS = state-trait anger scale; OSAB = observation scale for aggressive behavior.

* $p < .05$; ** $p < .01$ (two-tailed).

$p < .01$. The multiple correlation coefficient was .29, indicating that approximately 8.4% of the variance of irritation could be accounted for by the antisocial facet scores. The other facets of the PCL-R were not entered into the equation (all p 's $> .33$). Aggressive behavior was only significantly related to the lifestyle facet in the group of personality-disordered patients, $F(1, 157) = 12.71, p < .01$. The multiple correlation coefficient was .27, indicating that approximately 7.5% of the variance of aggressive behavior could be accounted for by the lifestyle facet scores. The other facets of the PCL-R were not entered into the equation (all p 's $> .62$). None of the facets of the PCL-R was significantly related to prosocial behavior in the group of personality-disordered patients. In the group of chronically psychotic patients, results indicated that the lifestyle facet of the PCL-R was significantly related to irritation, $F(1, 92) = 5.89, p < .02$. The multiple correlation coefficient was .25, indicating that approximately 6.0% of the variance of irritation could be accounted for by the lifestyle facet score. The other facets were not entered into the equation (all p 's $> .24$). For the group of psychotic patients, none of the facets of the PCL-R was significantly related to aggressive behavior or prosocial behavior (Table 5).

Change in behavior on the ward over a period of three years

The change of behavior was studied in the 115 patients (70 personality-disordered and 45 chronically psychotic patients) with seven measurements. The 45 patients who also stayed three years or more in the hospital but from whom less than five data measurements could be collected (Table 2) did not significantly differ from the 115 patients with seven measurements in PCL-R total score. Therefore, we assume that the studied group of 115 patients was representative of all patients who stayed three years or longer in the hospital.

For the scores on the Irritation/Anger subscale of the OSAB, both the assumptions of homogeneity of variance matrices ((Box's $M = 33.58, F(28, 30926.168) = 1.25, p = .17$)) and sphericity (Greenhouse Geisser $\epsilon = .87$) were met. The effect of the measurement within-subject factor Irritation/Anger was significant ((Wilks' Lambda $F(6, 108) = 5.107, p = < .001$)) and the effect size was large (partial $\eta^2 = .221$). Both the between-subject factor group (($F(1, 113) = 4.682, p = .033$, partial $\eta^2 = .040$)) and the interaction effect group \times irritation/anger measurement ((Wilks' Lambda $F(6, 108) = 2.031, p = .068$, partial $\eta^2 = .101$)) were not significant. A polynomial contrast analysis showed a significant quadratic trend effect, $F(1, 113) = 16.402, p = < .001$, partial $\eta^2 = .127$. The observed averages of irritation/anger on each time in Figure 1 per group show that patients on average became more irritated/angry during the first measurements, then their irritation dropped and slightly increased on the last measurement.

Regarding the scores on the OSAB subscale Aggressive Behavior, both the assumptions of homogeneity of variance matrices ((Box's $M = 38.04,$

Table 5. Stepwise multiple regression analyses between facets of PCL-R, and OSAB subscales for personality-disordered ($n = 159$) and chronically psychotic patients ($n = 94$), assessed shortly after admission (measurement 1).

OSAB Subscale	Personality-disordered patients					Chronically psychotic patients						
	PCL-R	<i>B</i>	<i>SE B</i>	95% CI	<i>p</i>	<i>R</i> ²	PCL-R	<i>B</i>	<i>SE B</i>	95% CI	<i>p</i>	<i>R</i> ²
Irritation/anger	Facet 4	0.354	0.093	[0.17–0.54]	.00	.084	Facet 3	0.32	0.13	[0.06–0.59]	.02	.060
Aggressive behavior	Facet 4	0.493	0.138	[0.22–0.77]	.00	.075	–	–	–	–	–	–
Prosocial behavior	–	–	–	–	–	–	–	–	–	–	–	–

Note: PCL-R = psychopathy checklist-revised; OSAB = observation scale for aggressive behavior.




Figure 1. Course of behavior during the first three years of stay for 70 personality-disordered and 45 chronically psychotic patients, measured with the OSAB subscales Irritation/anger, aggressive behavior, and prosocial behavior.

Note: Irritation/anger scores run from 9 to 36, Aggressive behavior scores from 10 to 40, and Prosocial scores from 12 to 48.

$F(28, 30926.168) = 1.263, p = .16$) and sphericity (Greenhouse Geisser $\epsilon = .81$) were also met. None of the effects were significant (within-subject factor: Wilks' Lambda $F(6, 108) = 1.056, p = .394$, partial $\eta^2 = .055$; between-subject factor: $F(1, 113) = 0.25, p = .620$, partial $\eta^2 = .002$; interaction aggressive behavior \times group: Wilks' Lambda $F(6, 108) = .988, p = .437$, partial $\eta^2 = .052$). Figure 1 shows the observed average aggressive behavior level per group.

For the scores on the OSAB subscale Prosocial Behavior, the assumption of homogeneity of variance matrices was violated, Box's $M = 50.63, F(28, 30926.168) = 1.680, p = .014$. Levene's test of equality of variances shows that the variances on measurement 2, 3, 5, and 7 were unequal, with more variance in the chronically psychotic group than in the personality-disordered group. Because the psychotic group is much smaller than the personality-disordered group, the F -test of the between-subject factor group and the interaction groups \times prosocial behavior is liberal (Stevens, 2007, p. 58). We, therefore, decided to use an alpha level of .01. The sphericity assumption was just about met (Greenhouse Geisser $\epsilon = .76$). The effect of the within-subject factor Irritation measurement was significant ((Wilks' Lambda $F(6, 108) = 4.352, p = .001$)), but the effect size was small (partial $\eta^2 = .195$). The between-subject factor was also significant, $F(1, 113) = 20.80, p < .001$, partial $\eta^2 = .155$. On average, the personality-disordered group scored higher on prosocial behavior than the chronically psychotic group. The interaction effect group \times prosocial behavior measurement ((Wilks' Lambda $F(6, 108) = 0.864,$

$p = .52$, partial $\eta^2 = .046$) was not significant. A polynomial contrast analysis showed a significant quadratic trend effect, $F(1, 113) = 8.69$, $p = .004$, partial $\eta^2 = .710$. The observed averages of prosocial behavior at each time point in Figure 1 show that patients on average showed more prosocial behavior during the first measurements, that their prosocial behavior then dropped and increased slightly again on the last measurement.

Because of the decreasing number of patients with an increasing number of measurement moments (see Table 1) and in order to corroborate our findings with the mixed analyses of variance, a comparison was performed on the OSAB subscales Irritation/Anger, Aggressive Behavior, and Prosocial Behavior with paired t -tests between measurement 2 and measurement 1 ($n = 248$), between measurement 3 and measurement 1 ($n = 236$), and so on to measurement 7 and measurement 1 ($n = 115$, see also Table 1). Similar results were found with these t -tests as with the mixed analyses of variance.

Predictors of behavior change after three years

During measurement 7, no relationships were found between PCL-R and OSAB scores in the personality-disordered group, but in the chronically psychotic group both irritation/anger ($r = .38$, $p < .05$) and prosocial behavior ($r = .33$, $p < .05$) turned out to be related with the Lifestyle facet of the PCL-R.

When we correlated residual change scores (measurement 7 versus measurement 1) on the three OSAB subscales with PCL-R, NEO-FFI, and STAS, we found that higher scores on the PCL-R facet Antisocial in the personality-disordered patients indicated a larger increase in prosocial behavior ($r = .29$, $p < .05$). In the chronically psychotic patients, no relationships were found between PCL-R facets and behavior change.

Discussion

Behavior on the wards of a forensic psychiatric institution was studied in a group of personality-disordered patients and a group of chronically psychotic patients. In accordance with the findings of Walters and Heilbrun (2010), in the personality-disordered patients, institutional aggression shortly after admission turned out to be related to the antisocial facet of psychopathy. However, in the chronically psychotic group, a relationship was found only between irritation/anger and the psychopathy lifestyle facet. After three years, no relationships between institutional aggression and any PCL-R facet were found in the personality-disordered group. In the chronically psychotic group, however, the psychopathy lifestyle facet again turned out to be related to irritation/anger and this point in time also to prosocial behavior. That no relationship between aggression on the ward and any PCL-R facet was found in the personality-disordered patients after a stay of three years in hospital supports the findings

of Heilbrun et al. (1998). In a group of 218 mentally disordered offenders, these researchers found significant correlations between aggressive incidents and PCL-R scores during the first two months of their stay in hospital, but these associations were no longer significant during the last two months of hospitalization. Regarding the different findings in the chronically psychotic group, it should be noted that both patient groups stayed in a similar controlled and structured environment. Probably, the personality-disordered patients in this study are more similar to North American prison samples and the chronically psychotic patients more similar to North American forensic psychiatric samples with especially schizophrenic or other psychotically disordered patients (Hildebrand & De Ruiter, 2012).

In both groups, irritation/anger and aggression on the ward were already low shortly after admission and did not decrease further during their stay of three years in hospital. However, prosocial behavior increased significantly in the two groups during this period. In the personality-disordered patients, the psychopathy antisocial facet was found to be related to a change in prosocial behavior, whereas none of the PCL-R facets was related to irritation/anger, aggressive behavior, or prosocial behavior in the chronically psychotic group.

Our study had a number of limitations. First, PCL-R scores were based on file study alone and most scores were single rated. In a sample of 41 inpatients, the intra-class correlation coefficient of the PCL-R total score was excellent, but the coefficients of the four facets varied from fair to excellent. Second, in 8% of the cases that were studied on behavior change during a three-year period, we had to interpolate data for the observation scale because of missing values. Third, some patients refused to complete the self-report questionnaires, and the patients who responded might have done this in a more or less socially desirable way. Consequently, the number of assessed patients was rather low for some measures. Fourth, the validity of the NEO-FFI and STAS has not been studied in Dutch forensic psychiatric patients until now. Finally, with the observation scale, no difference could be made between reactive and proactive aggression. However, because most aggressive behavior was exhibited in interactions with the staff, we suppose that this aggressive behavior was mostly reactive of sorts. This would explain why no relationships were found between the psychopathy interpersonal and affective facets as in the Laurell, Belfrage, and Hellström (2010) study.

In spite of these limitations, several provisional conclusions can be drawn. For example, the structured and controlled environment of the forensic psychiatric hospital probably had an attenuating effect on the aggressive behavior of most patients. This effect is in line with earlier studies, in which inpatients turned out to score lower than forensic psychiatric outpatients on multiple-choice self-report questionnaires measuring anger and aggression (Hornsveld, Muris, & Kraaimaat, 2011; Hornsveld, Muris, Kraaimaat, & Meesters, 2009). Vitacco et al. (2009) explained the various results of the studies on institutional behavior through the insufficient uniformity in definitions of institutional

aggression and the lack of differentiation between reactive and proactive aggression. They also noticed that base rates of institutional aggression are generally low, due to medication and the presence of staff.

Our findings and those of others may illustrate the problems by demonstrating the effects of treatment programs for inpatients when the outcome is based only on negative behavior, such as hostility, anger, or aggression. It is, therefore, recommended to study positive, socially desirable behavior additionally. A recent study by De Vries Robbé, De Vogel, and Douglas (2013) indicates that protective factors, which mostly refer to positive behavior, may contribute to a more valid assessment of recidivism risk.

In our opinion, the present study may indicate that for a considerable group of patients, a three- to four-year stay in a Dutch forensic psychiatric institution should suffice for assessment of risk and needs, establishing a treatment plan, and executing this plan through treatment programs. Therefore, with an average stay of 9.8 years for Dutch forensic psychiatric inpatients in mind, it is probably worthwhile to investigate whether this period can be shortened by transferring these patients to a less-structured environment, in order to safely practice their newly acquired behavior. Such an environment would also allow professionals to monitor the patients' behavior, assess treatment results, and intervene in case of an unexpected relapse. Further research may indicate if a policy focusing on a safe but more gradual return of those inpatients in society is cheaper, and also increasing treatment efficacy along with.

References

- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders (DSM-IV-TR)*. Washington, DC: Author.
- Baker, R.D., & Hall, J.N. (1988). REHAB: A new assessment instrument for chronic psychiatric patients. *Schizophrenia Bulletin*, *14*, 97–111. doi:10.1093/schbul/14.1.97
- Belfrage, H., & Douglas, K.S. (2002). Treatment effects on forensic psychiatric patients measured with the HCR-20 violence risk assessment scheme. *International Journal of Forensic Mental Health*, *1*, 25–36. doi:10.1080/14999013.2002.10471158
- Chakhssi, F., De Ruiter, C., & Bernstein, D. (2010). Change during forensic treatment in psychopathic versus nonpsychopathic offenders. *Journal of Forensic Psychiatry and Psychology*, *21*, 660–682. doi:10.1080/14789949.2010.483283
- Cicchetti, D.V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, *6*, 284–290.
- Cornell, D.G., Peterson, C.S., & Richards, H. (1999). Anger as a predictor of aggression among incarcerated adolescents. *Journal of Consulting and Clinical Psychology*, *67*, 108–115. doi:10.1037/0022-006X.67.1.108
- Costa, P.T., & McCrae, R.R. (1989). *NEO personality inventory-revised*. Port Huron, MI: Sigma Assessment Systems.
- Costa Jr, P.T., & McCrae, R.R. (1992). *Revised NEO personality inventory (NEO-PI-R) and the five factor inventory (NEO-FFI): Professional manual*. Odessa, FL: Psychological Assessment Resources.

- De Beurs, E., & Barendregt, M. (2008). *Mogelijkheden voor therapie-effectonderzoek in de tbs-sector: komen tot een evidence base onder zorgprogramma's* [Possibilities for therapy effect-research in the tbs-sector: Arriving at an evidence base for treatment programs]. Utrecht/The Hague, NL: NIFP/WODC.
- De Jonge, E., Nijman, H.L.I., & Lammers, S.M.M. (2009). Gedragsverandering tijdens tbs-behandeling: een multicenter onderzoek [Behavior change during tbs-treatment: A multi-center study]. *Tijdschrift voor Psychiatrie* [Dutch Journal of Psychiatry] *51*, 205–215.
- De Vries Robbé, M., De Vogel, V., & Douglas, K.S. (2013). Risk factors and protective factors: A two-sided dynamic approach to violence risk assessment. *The Journal of Forensic Psychiatry and Psychology*, *24*, 440–457. doi:10.1080/14789949.2013.818162
- Edens, J.F., Campbell, J.S., & Weir, J.M. (2007). Youth psychopathy and criminal recidivism: A meta-analysis of the psychopathy checklist measures. *Law and Human Behavior*, *31*, 53–75. doi:10.1007/s10979-006-9019-y
- Guy, L.S., Edens, J.F., Anthony, C., & Douglas, K.S. (2005). Does psychopathy predict institutional misconduct among adults: A meta-analytic investigation. *Journal of Consulting and Clinical Psychology*, *73*, 1056–1064. doi:10.1037/0022-006X.73.6.1056
- Hare, R.D. (1980). The assessment of psychopathy in criminal populations. *Personality and Individual Differences*, *1*, 111–119. doi:10.1016/0191-8869(80)90028-8
- Hare, R.D. (1991). *The Hare Psychopathy Checklist-Revised*. Toronto: Multi-Health Systems.
- Hare, R.D. (2003). *The hare psychopathy checklist-revised*. (2nd ed.). Toronto: Multi-Health Systems.
- Hare, R.D., & Neumann, C.S. (2006). The PCL-R assessment of psychopathy: Development, structural properties, and new directions. In: C.J. Patrick (Ed.), *Handbook of Psychopathy* (pp. 58–88). New York, NY: Guilford Press.
- Hart, S.D., Cox, D.N., & Hare, R.D. (1995). *Hare psychopathy checklist: Screening version*. Toronto: Multi-Health Systems.
- Heilbrun, K., Hart, S.D., Hare, R.D., Gustafson, D., Nunez, C., & White, A.J. (1998). Inpatient and postdischarge aggression in mentally disordered offenders: The role of psychopathy. *Journal of Interpersonal Violence*, *13*, 514–527. doi:10.1177/088626098013004007
- Hildebrand, M., & De Ruiter, C. (2012). Psychopathic traits and change on indicators of dynamic risk factors during inpatient forensic psychiatric treatment. *International Journal of Law and Psychiatry*, *35*, 276–288. doi:10.1017/j.ijlp.2012.04.001
- Hildebrand, M., De Ruiter, C., & Nijman, H. (2004). PCL-R psychopathy predicts disruptive behavior among male offenders in a Dutch forensic psychiatric hospital. *Journal of Interpersonal Violence*, *19*, 13–29. doi:10.1177/0886260503259047
- Hoekstra, H.A., Ormel, J., & De Fruyt, F. (1996). *Handleiding NEO Persoonlijkheidsvragenlijsten* [Manual NEO personality questionnaires]. Lisse: Swets Test Services.
- Hornsveld, R.H.J., Muris, P.E.H.M., & Kraaimaat, F.W. (2011). The Novaco anger scale-provocation inventory (1994 version) in Dutch forensic psychiatric patients. *Psychological Assessment*, *23*, 937–944. doi:10.1037/a0024018
- Hornsveld, R.H.J., Muris, P., Kraaimaat, F.W., & Meesters, C. (2009). Psychometric properties of the aggression questionnaire in Dutch violent forensic psychiatric patients and secondary vocational students. *Assessment*, *16*, 181–192. doi:10.1177/1073191108325894

- Hornsveld, R.H.J., Nijman, H.L.I., Hollin, C.R., & Kraaiaat, F.W. (2007). Development of the observation scale for aggressive behavior (OSAB) for Dutch forensic psychiatric inpatients with an antisocial personality disorder. *International Journal of Law and Psychiatry*, 30, 480–491. doi:10.1016/j.ijlp.2007.09.009
- Hornsveld, R.H.J., Nijman, H.L.I., & Kraaiaat, F.W. (2008). Aggression Control Therapy for violent forensic psychiatric patients: First results. *Psychology, Crime and Law*, 14, 1–18.
- Kennealy, P.J., Skeem, J.L., Walters, G.D., & Camp, J. (2010). Do core interpersonal and affective traits of the PCL-R psychopathy interact with antisocial behavior and disinhibition to predict violence? *Psychological Assessment*, 22, 569–580. doi:10.1037/a0019618
- Kerlinger, F. (1975). *Foundations of behavioral research*. New York, NY: Holt, Rinehart, and Winston.
- Laurell, J., Belfrage, H., & Hellström, Å. (2010). Facets on the psychopathy checklist screening version and instrumental violence in forensic psychiatric patients. *Criminal Behaviour and Mental Health*, 20, 285–294. doi:10.1002/cbm.779
- Leistico, A.R., Salekin, R.T., DeCoster, J., & Rogers, R. (2008). A large-scale meta-analysis relating the Hare measures of psychopathy to antisocial conduct. *Law and Human Behavior*, 32, 28–45. doi:10.1007/s10979-007-9096-6
- McGuire, J. (2013). ‘What works’ to reduce re-offending. 18 Years on. In: L. A. Craig, L. Dixon, & T. A. Gannon (Eds.), *What works in offender rehabilitation: An evidence-based approach to assessment and treatment* (pp. 20–49). Chichester: Wiley. doi: 10.1002/9781118320655
- Ministry of Security and Justice. (2002). *Handleiding HKT-30. Risicotaxatie in de Forensische Psychiatrie* [Manual HKT-30. Risk assessment in forensic psychiatry]. The Hague: Ministry of Safety and Justice.
- Nachtegaal, M.H., Van der Horst, R.P., & Schönberger, H.J.M. (2011). *Inzicht in de verblijfsduur van tbs-gestelden* [Insight in the length of stay for offenders detained by hospital order]. Den Haag: Boom Juridische uitgevers.
- Nijman, H., De Kruijk, C., & Van Nieuwenhuizen, Ch. (2004). Behavioral changes during forensic psychiatric (TBS) treatment in the Netherlands. *International Journal of Law and Psychiatry*, 27, 79–85. doi:10.1016/j.ijlp.2003.12.001
- Reed, V., Woods, P., & Robinson, D. (2000). *Behavioural status index (BEST-Index): A ‘life skills’ assessment for selecting and monitoring therapy in mental health care*. Sheffield: Psychometric Press.
- Rorschach, H. (1942). *Psychodiagnostics: A diagnostic test based on perception*. Bern: Hans Huber (Original work published 1921).
- Singh, J.P., Grann, M., & Fazel, S. (2011). A comparative study of violence risk assessment tools: A systematic review and metaregression analysis of 68 studies involving 25,980 participants. *Clinical Psychology Review*, 31, 499–513. doi:10.1016/j.cpr.2010.11.009
- Sloore, H., Derksen, J.J.L., Hellenbosch, G., & De Mey, H.R.A. (1993). *De Minnesota Multiphasic Personality Inventory-II (Dutch translation)*. Nijmegen, NL: PEN Test Publishers.
- Spielberger, C.D. (1980). *Preliminary manual for the state-trait anger scale (STAS)*. Tampa: University of South Florida.
- Spielberger, C.D. (1988). *State-trait anger expression inventory: Professional manual*. (research ed.). Odessa: Psychological Assessment Resources.
- Stevens, J.P. (2007). *Intermediate statistics: A modern approach* (3rd ed.). New York, NY: Taylor & Francis.

- Timmerman, I.G.H., Vastenburg, N.C., & Emmelkamp, P.M.G. (2001). The forensic inpatient observation scale (FIOS): Development and validity. *Criminal Behavior and Mental Health, 11*, 144–162.
- Van der Ploeg, H.M., Defares, P.B., & Spielberger, C.D. (1982). *Handleiding bij de Zelf-Analyse Vragenlijst, ZAV* [Manual for the Self-Analysis Questionnaire, ZAV]. Lisse: Swets & Zeitlinger.
- Van Marle, H.J.C. (2002). The Dutch entrustment act (TBS): It's principles and innovations. *International Journal of Forensic Mental Health, 1*, 83–92. doi:10.1080/14999013.2002.10471163.
- Vertommen, H., Verheul, R., De Ruiter, C., & Hildebrand, M. (2002). *De herziene versie van Hare's Psychopathie Checklist (PCL-R)* [The revised version of Hare's Psychopathy Checklist (PCL-R)]. Lisse: Swets Test Publishers.
- Vitacco, M.J., Van Rybroek, G.J., Rogstad, J.E., Yahr, L.E., Tomony, J.D., & Saewert, E. (2009). Predicting short-term institutional aggression in forensic patients: A multi-trait method for understanding subtypes of aggression. *Law and Human Behavior, 33*, 308–319. doi:10.1007/s10979-008-9155-7
- Walters, G.D., & Heilbrun, K. (2010). Violence risk assessment and facet 4 of the psychopathy checklist: Predicting institutional and community aggression in two forensic samples. *Assessment, 17*, 259–268. doi:10.1177/1073191109356685
- Walters, G.D., Knight, R.A., Grann, M., & Dahle, K. (2008). Incremental validity of the psychopathy checklist facet scores: Predicting release outcome in six samples. *Journal of Abnormal Psychology, 117*, 396–405. doi:10.1037/0021-843X.117.2.396
- Webster, C.D., Douglas, K.S., Eaves, D., & Hart, S.D. (1997). *HCR-20: Assessing risk for violence (version 2)*. Burnaby: Mental Health, Law, and Policy Institute, Simon Fraser University.
- Yang, M., Wong, S.C.P., & Coid, J. (2010). The efficacy of violence prediction: A meta-analytic comparison of nine risk assessment tools. *Psychological Bulletin, 136*, 740–767. doi:10.1037/a0020473