

Psychological characteristics of Dutch violent forensic psychiatric in- and outpatients

Ruud H.J. Hornsveld (Ph.D.),
clinical psychologist/researcher,
Erasmus University Medical Center

r.hornsveld@tiscali.nl

www.Agressiehanteringstherapie.nl

Contents

Part 1

- **“Models” for violent behavior**
- **New measurement instruments**
- **Psychological characteristics**
- **Aggression Replacement Training**

Part 2

- **Course of antisocial and prosocial behavior during the first three years of stay**
- **Predictors of behavior on the ward**
- **Consequences for treatment programs**

“Models” for violent behavior

Risk-Need-Responsivity “Model” (Andrews & Bonta, 2010)

Criminogenic needs (“Central Eight”):

- **history of antisocial behavior**
- **antisocial personality pattern**
- **antisocial cognitions**
- **antisocial associates**
- **family/marital circumstances**
- **school/work**
- **leisure/recreation**
- **substance abuse**

Good Lives Model (Ward & Marshall, 2004)


Ten good live goals;

- healthy life
- knowledge
- excellence in play and work
- agency
- inner peace
- relatedness
- community
- spirituality
- happiness
- creativity

Comments on Risk-Need-Responsivity model and Good Lives Model

- **Items have different abstraction levels**
- **Not that so much models, but more a list of areas for special attention**
- **Translation of each area in psychological characteristics is needed: functional analyses**

Model for aggressive behavior



New measurement instruments

Comments on most measurement instruments, used in forensic psychiatry

Problems

- **Self-report questionnaires mostly developed using college or university students**
- **No investigation on the psychometric properties of these questionnaires in Dutch forensic psychiatric in- and outpatients**
- **Observation scales about aggression mostly developed for general psychiatric patients in crisis**

Findings

- **Questionnaires are reliable and valid when used for Dutch forensic psychiatric patients, but most divisions in subscales do not fit**

Current measurement instruments for program evaluation

Personality traits

- ***Psychopathy Checklist-Revised (PCL-R;*** Vertommen, Verheul, De Ruiter, & Hildebrand, 2002): psychopathy
- ***NEO Five-Factor Inventory (NEO-FFI;*** Hoekstra, Ormel, & De Fruyt, 1996): neuroticism, agreeableness, and conscientiousness
- **Trait Anger subscale of the Spielberger (1980) *State-Trait Anger Scale (STAS;*** Van der Ploeg, Defares, & Spielberger, 1982): general disposition to anger

Current measurement instruments for program evaluation (continued)

Behavior

- ***Aggression Questionnaire (AQ; Hornsveld, Muris, Kraaimaat, & Meesters, 2009): general aggression and physical aggression***
- ***NAS part of the Novaco Anger Scale-Provocation Inventory (NAS-PI; Hornsveld, Muris, & Kraaimaat, 2011): anger***
- ***Inventory of Interpersonal Situations (IIS; Van Dam-Baggen & Kraaimaat, 1999): social anxiety and frequency of social skills***

New instruments for program evaluation

- **Adapted version of Rosenzweig Picture-Frustration Study (PFS-AV: Hornsveld, Nijman, Hollin, & Kraaimaat, 2007a): hostility**
- **Observation Scale for Aggressive Behavior (OSAB; Hornsveld, Nijman, Hollin, & Kraaimaat, 2007b): behavior of forensic psychiatric inpatients on the ward**

Adapted Version of Rosenzweig Picture-Frustration Study (PFS-AV): Hostility



You are not allowed to cross the street when the light is red.

Observation Scale for Aggressive Behavior (OSAB; Hornsveld et al., 2007)

Six subscales:

- **Irritation/anger (5 items)**
- **Anxiety/Gloominess (4 items)**
- **Aggressive behavior (10 items)**
- **Prosocial behavior (12 items)**
- **Antecedents (6 items)**
- **Sanctions (3 items)**

Scoring: Behavior on the ward during last week

Scores of subscales


Scoring of items: 'no' = 1, 'seldom' = 2, 'occasionally' = 3, and 'frequently' = 4

Range of subscale scores:


- **Irritation/anger: 5 - 20**
- **Anxiety/Gloominess: 4 - 16**
- **Aggressive behavior: 10 - 40**
- **Prosocial behavior: 12 - 48**
- **Antecedents: 6 - 24**
- **Sanctions: 3 -13**

Dynamic criminogenic needs of forensic psychiatric patients


Inpatients versus norms ($N = 136$)




Inpatients versus norms ($N = 136$)




Inpatients ($N = 136$) versus detainees ($N = 100$)




Inpatients ($N = 136$) versus detainees ($N = 100$)




Outpatients versus norms (N = 200)



Outpatients versus norms (N = 200)



Young outpatients ($N = 123$) versus male students ($N = 160$)



Aggression Replacement Training (ART)

Participants

Data sets

- 123 outpatients (mean age = 17.35 years, $SD = 1.82$, range: 15-21 years)
- 73 patients were measured both during the intake interview and at the start of the training (mean age = 17.12 years, $SD = 1.72$, range: 15-21 years)
- 62 patients completed the questionnaires at both the start and the end of the training (mean age = 17.35 years, $SD = 1.91$, range: 15-21 years)
- 61 patients withdrew prematurely during the waiting period or during the training (nonstarters plus non-completers; mean age 17.35 years, $SD = 1.82$, range: 15-21 years)

Measures

- ***Psychopathy Checklist-Revised (PCL-R;***
Vertommen, Verheul, De Ruiter, & Hildebrand, 2002)
- ***NEO Five-Factor Inventory (NEO-FFI; Hoekstra,***
Ormel, & De Fruyt, 1996)
- **Trait Anger subscale of the Spielberger (1980)**
State-Trait Anger Scale (STAS; Van der Ploeg,
Defares, & Spielberger, 1982)
- ***Adapted Version of the Picture-Frustration Study***
(PFS-AV; Hornsveld, Nijman, Hollin, & Kraaimaat,
2007)
- ***Aggression Questionnaire (AQ; Hornsveld, Muris,***
Kraaimaat, & Meesters, 2009)

Measures (continued)

- **NAS** part of the *Novaco Anger Scale-Provocation Inventory* (NAS-PI; Hornsveld, Muris, & Kraaimaat, 2011)
- *Inventory of Interpersonal Situations* (IIS; Van Dam-Baggen & Kraaimaat, 1999)

Outpatient ART and design

Outpatient ART

Fifteen weekly sessions lasting 1½ hours each and three five-weekly follow-up meetings for six to eight patients:

- **anger management, sessions 1 to 5**
- **social skills, sessions 6 to 10**
- **moral reasoning, sessions 11 to 15**
- **follow-up and evaluation, sessions 16 to 18**

Participants had to complete homework assignments

Design

Three measurement moments:

- **at intake/before a waiting period**
- **after the waiting period/before the training**
- **after the training (post-training measurement)**

Results

Criminogenic needs

- Compared with a *reference group* of 275 secondary vocational students, patients scored higher on trait anger, hostility, and aggression, and lower on social anxiety

Nonstarters and noncompleters

- Patients who withdrew prematurely scored higher on psychopathy than the completers, in particular on the factor antisocial behavior

Intake measurement vs. pre measurement (*n* = 73)

Measure	Content of scale	<i>M</i> (<i>SD</i>)		Effect <i>d</i>
		Intake	Pre	
PFS-AV	Hostility	33.22 (9.58)	34.16 (11.49)	-.13
AQ	Aggression	90.00 (27.88)	85.59 (21.57)	.21
	Phys. aggr.	33.01 (18.47)	29.48 (8.19)	.36
NAS-PI	Anger	87.52 (17.35)*	90.81 (19.32)*	-.29
IIS	Social anxiety	71.43 (28.73)	68.07 (25.80)	.24
	Social skills	112.42 (25.19)	112.32 (25.18)	.01

* $p < .05$

Pre measurement vs. post measurement ($n = 62$)

Measure	Content of scale	<i>M (SD)</i>		Effect <i>d</i>
		Intake	Pre	
PFS-AV	Hostility	33.34 (12.30)	30.84 (12.27)	.25
AQ	Aggression	82.56 (20.67)	78.90 (20.32)	.21
	Phys. aggr.	28.39 (8.02)*	26.45 (7.46)*	.28
NAS-PI	Anger	87.29 (18.31)	83.98 (16.74)	.21
IIS	Social anxiety	65.36 (22.75)*	57.74 (22.75)*	.31
	Social skills	115.88 (22.22)	116.93 (29.75)	-.04

* $p < .05$

Results (continued)

Behavior change

- **No change in 73 patients between intake and pre measurement, except for an increase in anger**
- **Compared with the pre training measurement, 62 patients scored lower on physical aggression and social anxiety during the post-training measurement. There was a trend in the reduction of hostility, aggression, and anger**
- **After completion of the training, patients did not differ from the *reference group* of secondary vocational students with respect to hostility and aggressive behavior**

Discussion

Drop-out

- **In the current study, 61 of the 123 patients did not show up at the start of the training or did not complete the training**
- **This result is in line with the results of other studies on treatment dropouts (e.g., Olver & Wong, 2009)**
- **Non-completion has been associated with a higher risk of recidivism (Wormith, Olver, Stevenson, & Girard, 2007), as well as aggression and rule-violating behaviors (Beyko & Wong, 2005)**
- **There seems to be a relation between psychopathy, treatment attrition, and recidivism risk**

Consequences for treatment

- **For this group of patients a more consequent and stricter policy is required among the referring agencies in case of drop-out**
- **Refusing to follow the training hardly had any negative consequences in most cases**
- **Creating alternative conditions and consequences for the completion of an obligatory treatment program has the highest priority**
- **For instance, the training can be provided at the office of the after-care and resettlement organization by a qualified trainer from the outpatient clinic and a probation officer**

Behavior of Dutch violent forensic psychiatric inpatients on the ward

**Ruud H.J. Hornsveld (Ph.D.),
clinical psychologist/researcher,
Erasmus University Medical Center**

r.hornsveld@tiscali.nl

www.Agressiehanteringstherapie.nl

Observation Scale for Aggressive Behavior (OSAB; Hornsveld et al., 2007)

Three of the six subscales:

- **Irritation/anger (5 items)**
- **Aggressive behavior (10 items)**
- **Prosocial behavior (12 items)**

Scoring: Behavior on the ward during last week

Table 1. Number of patients assessed on seven measurements

	Total group		Personality disordered		Chronically psychotic	
	<i>N</i>	<i>M (SD)</i>	<i>N</i>	<i>M(SD)</i>	<i>N</i>	<i>M(SD)</i>
1	253	37.49 (10.38)	159	38.62 (10.64)	94	35.56 (9.69)
2	248	37.40 (10.44)	157	38.58 (10.68)	91	35.37 (9.75)
3	236	37.17 (10.41)	148	38.38 (10.66)	88	35.15 (9.69)
4	213	37.20 (10.34)	134	38.69 (10.88)	79	34.67 (8.86)
5	178	36.97 (10.46)	108	38.69 (11.14)	70	34.30 (8.74)
6	146	37.03 (10.17)	86	38.99 (10.77)	60	34.22 (8.58)
7	115	36.97 (10.27)	70	38.79 (10.75)	45	34.13 (8.85)

Figure 1. Course of irritation/anger and aggressive behavior during the first three years of stay




Figure 2. Course of prosocial behavior during the first three years of stay




Figure 3. Course of irritation/anger and aggressive behavior during the first three years of stay




Figure 4. Course of prosocial behavior during the first three years of stay




Table 2. Mean scores and SD's in personality disordered and chronically psychotic patients

Meas- ure	Factor or sub- scale	Personality disordered		Chronically psychotic		Differences between subsamples
		<i>N</i>	<i>M (SD)</i>	<i>N</i>	<i>M (SD)</i>	
PCL-R	Psychop	159	22.25 (8.06)	94	17.96 (7.84)	$F(2,250) = 13.13 (p < .001)^{**}$
	Interpers	159	3.57 (2.47)	94	1.80 (1.88)	$F(2,250) = 18.02 (p < .001)^{**}$
	Affective	159	6.13 (1.72)	94	5.64 (1.84)	$F(2,250) = 3.10 (p = .047)^*$
	Lifestyle	159	5.94 (2.76)	94	4.98 (2.90)	$F(2,250) = 14.01 (p < .001)^{**}$
	Antisoc	159	5.11 (2.81)	94	4.68 (2.81)	$F(2,250) = 6.58 (p = .002)^{**}$
NEO-FFI	Neurot	97	32.24 (8.49)	48	31.73 (7.92)	$F(2,142) = 0.95 (p = .909)$
	Agree	97	41.59 (5.23)	48	42.52 (4.93)	$F(2,142) = 0.53 (p = .591)$
STAS	Anger	92	17.91 (6.64)	47	15.85 (4.29)	$F(2,136) = 1.94 (p = .148)$

Table 4. Correlations assessed shortly after admittance

Measure	Factors or subscales	Personality disordered patients			Chronically psychotic patients		
		Irritation/ Anger	Aggressive behavior	Pro-social behavior	Irritation/ anger	Aggressive behavior	Pro-social behavior
PCL-R	Psychp	.236**	.208**	.016	.169	.052	.080
	Interper	.097	.069	.057	.192	.056	.089
	Affect	.177*	.122	-.051	.078	-.012	.073
	Lifest	.207**	.199*	-.001	.245*	.160	.034
	Antisoc	.290**	.274**	.091	.051	-.030	.039
NEO-FFI	Neurot	.199*	.209*	-.107	.059	-.001	.161
	Agree	-.097	-.160	.093	-.111	-.127	-.099
STAS	Anger	.140	.214*	.023	.157	.182	.078

**Table 5. Change scores on the OSAB
(measurement 1 vs. measurement 7)**

OSAB sub- scale	Personality disordered patients			Chronically psychotic patients		
	Measure- ment 1	Measure- ment 7	Ef- fect size	Measure- ment 1	Measure- ment 7	Ef- fect size
	M (SD)	M (SD)	d	M (SD)	M (SD)	d
Irrit./Anger	10.57 (3.31)	10.46 (3.04)	.057	9.60 (4.03)	9.36 (3.57)	.113
Aggr. beh.	14.93 (5.38)	15.16 (4.67)	.065	15.02 (6.15)	15.07 (5.79)	.015
Prosoc. beh.	29.50 (8.39)	32.70 (6.94)	.650	24.22 (7.99)	27.38 (7.60)	.646

Table 6. *Outflow of patients*

Patients	Percentage	Age	PCL-R	Aggression on the ward
7 measurements	56.4	36.97 (10.27)	20.49 (7.97)	14.97 (5.66)
3 year of stay but no 7 measurements	22.1	39.38 (11.78)	17.68 (7.94)	15.92 (4.60)
Reselection	9.8	36.95 (8.89)	22.63 (7.86)	16.16 (5.23)
Long-stay	2.9	55.00 (8.46)	23.40 (7.57)	18.40 (2.61)
Finishing TBS or leave	3.4	41.29 (11.94)	17.00 (8.25)	14.14 (2.48)
Others	5.4	35.89 (6.31)	25.67 (9.35)	15.67 (3.20)

Conclusions

- **No relation between length of stay and mood, aggressive behavior, and sanctions.**
- **However, social skills are related to length of stay.**
- **Personality disordered patients exhibit more anger, more aggressive behavior, and more prosocial behavior than chronic psychotic patients.**
- **Patients with relatively high scores on the**
- **PCL-R exhibit more anger, more aggressive behavior, but also more prosocial behavior than patients with relatively low scores on the PCL-R.**
- **In general, base rates of negative behaviors are low.**

Discussion and recommendations

- **Outcome of treatment programs should not be based on negative but on positive behavior.**
- **Limited validity of risk assessment instruments if they are based on negative behaviors.**
- **Protective factors, which refer to positive behavior, can contribute considerably to a better prediction of recidivism risk (SAPROF).**
- **There is a group of inpatients for which a stay longer than three or four years has no incremental value.**

References

- Andrews, D. A., & Bonta, J. (2010). *The Psychology of Criminal Conduct* (fifth edition). Cincinnati, OH: Anderson.
- Goldstein, A. P., Glick, B., & Gibbs, J. C. (1998). *Aggression Replacement Training: A comprehensive intervention for aggressive youth*. Champaign, IL: Research Press.
- Hoekstra, H. A., Ormel, J., & De Fruyt, F. (1996). *Handleiding NEO Persoonlijkheidsvragenlijsten [Manual NEO Personality Questionnaires]*. Lisse, NL: Swets Test Services.
- Hornsveld, R. H. J., Kraaimaat, F. W., Bouwmeester, S., Polak, M. A., & Zwets, A. J. (2014). Behavior on the ward of personality-disordered inpatients and chronically psychotic inpatients during a three-year stay in a Dutch forensic psychiatric hospital. *The Journal of Forensic Psychiatry and Psychology*. doi: 10.1080/14789949.2014.947308
- Hornsveld, R. H. J., Kraaimaat, F. W., Muris, P., Zwets, A. J., & Kanters, T. (2014). Aggression Replacement Training for violent young men in a forensic psychiatric outpatient clinic. *Journal of Interpersonal Violence*. In press.

References (continued)

- Hornsveld, R. H. J., Muris, P. E. H. M., & Kraaimaat, F. W. (2011). The Novaco Anger Scale-Provocation Inventory (1994 version) in Dutch forensic psychiatric patients. *Psychological Assessment, 23*, 937-944. doi: 10.1037/a0024018
- Hornsveld, R. H. J., Muris, P., Kraaimaat, F. W., & Meesters, C. (2009). Psychometric properties of the Aggression Questionnaire in Dutch violent forensic psychiatric patients and secondary vocational students. *Assessment, 16*, 181-192. doi: 10.1177//1073191108325894
- Hornsveld, R. H. J., Nijman, H. L. I., Hollin, C. R., & Kraaimaat, F. W. (2007a). An adapted version of the Rosenzweig Picture-Frustration Study (PFS-AV) for the measurement of hostility in violent forensic psychiatric patients. *Criminal Behaviour and Mental Health, 17*, 45-56. doi: 10.1002/cbm.638
- Hornsveld, R. H. J., Nijman, H. L. I., Hollin, C. R., & Kraaimaat, F. W. (2007b). Development of the Observation Scale for Aggressive Behavior (OSAB) for Dutch forensic psychiatric inpatients with an antisocial personality disorder. *International Journal of Law and Psychiatry, 30*, 480-491. doi: 10.1016/j.ijlp.2007.09.009

References (continued)

- Van Dam-Baggen, C. M. J. & Kraaimaat, F. W. (1999). Assessing Social Anxiety: The Inventory of Interpersonal Situations (IIS). *European Journal of Psychological Assessment, 15*, 25-38.
doi: 10.1027//1015-5759.15.1.25
- Van der Ploeg, H. M., Defares, P. B., & Spielberger, C. D. (1982). *Handleiding bij de Zelf-Analyse Vragenlijst, ZAV [Manual for the Self-Analysis Questionnaire, ZAV]*. Lisse, NL: Swets & Zeitlinger.
- Vertommen, H., Verheul, R., De Ruiter, C., & Hildebrand, M. (2002). *De herziene versie van Hare's Psychopathie Checklist (PCL-R) [The revised version of Hare's Psychopathy Checklist (PCL-R)]*. Lisse, NL: Swets Test Publishers.
- Ward, T., & Marshall, W. L. (2004). Good lives, aetiology and the rehabilitation of sex offenders: A bridging theory. *Journal of Sexual Aggression: Special Issue: Treatment & Treatability, 10*, 153-169.
- Wormith, J. S., Olver, M. E., Stevenson, H. E., & Girard, L. (2007). The long-term prediction of offender recidivism using diagnostic, personality, and risk/need approaches to offender assessment. *Psychological Services, 4*, 287-305. doi: 10.1177/0093854802029004006